

SCIENTIFIC REPORTS

OPEN

Metformin stimulates IGFBP-2 gene expression through PPARalpha in diabetic states

Hye Suk Kang^{1,*}, Ho-Chan Cho^{2,*}, Jae-Ho Lee¹, Goo Taeg Oh³, Seung-Hoi Koo⁴,
Byung-Hyun Park⁵, In-Kyu Lee⁶, Hueng-Sik Choi⁷, Dae-Kyu Song¹ & Seung-Soon Im¹

Received: 11 October 2015

Accepted: 11 March 2016

Published: 24 March 2016

The anti-diabetic drug, metformin, exerts its action through AMP-activated protein kinase (AMPK), and Sirtuin (*Sirt1*) signaling. Insulin-like growth factor (IGF)-binding protein 2 (IGFBP-2) prevents IGF-1 binding to its receptors, thereby contributing to modulate insulin sensitivity. In this study, we demonstrate that metformin upregulates *Igfbp-2* expression through the AMPK-Sirt1-PPAR α cascade pathway. In the liver of high fat diet, *ob/ob*, and *db/db* mice, *Igfbp-2* expression was significantly decreased compared to the expression levels in the wild-type mice ($p < 0.05$). Upregulation of *Igfbp-2* expression by metformin administration was disrupted by gene silencing of *Ampk* and *Sirt1*, and this phenomenon was not observed in *Ppar α* -null mice. Notably, activation of IGF-1 receptor (IGF-1R)-dependent signaling by IGF-1 was inhibited by metformin. Finally, when compared to untreated type 2 diabetes patients, the metformin-treated diabetic patients showed increased IGFBP-2 levels with diminished serum IGF-1 levels. Taken together, these findings indicate that IGFBP-2 might be a new target of metformin action in diabetes and the metformin-AMPK-Sirt1-PPAR α -IGFBP-2 network may provide a novel pathway that could be applied to ameliorate metabolic syndromes by controlling IGF-1 bioavailability.

In biological fluids, the insulin-like growth factor binding protein (IGFBP) family proteins can complex with both IGF-I and II, and are regulators of IGF actions on metabolism and growth^{1,2}. Currently, there are six mammalian IGFBPs designated IGFBP-1–6 have been characterized³ and newly IGFBP-7 was identified as a member of the IGFBP superfamily⁴. The primary function of the IGFBPs is to restrict the bioavailability of IGF-1 in target tissues⁵. Among the IGFBPs, IGFBP-2 modulates IGF-1 bioactivity by interacting with IGF-1⁶. IGFBP-2 is highly expressed in the liver, adipocytes, and central nervous system, and is involved in metabolic homeostasis, insulin resistance, diabetes, and obesity^{7–9}. Moreover, IGFBP-2 is suggested to be used as a marker protein for metabolic dysfunction^{10,11}. IGFBP-2 is the abundant in blood, and has been shown to play a role in preventing insulin resistance and diet-associated obesity in mice⁹. However, the regulatory mechanism of IGFBP-2 expression and its clinical relevance to diabetic states in mice and humans remain poorly understood.

Sirtuin 1 (*Sirt1*), an NAD⁺-dependent protein deacetylase, is involved in controlling glucose, lipid homeostasis, aging, inflammation, and circadian regulation of metabolism and cellular processes¹². It regulates metabolic homeostasis by deacetylating crucial transcriptional factors such as peroxisome proliferator-activated receptor α (PPAR α), PPAR γ , farnesoid X receptor (FXR), liver X receptor α (LXR α), PPAR γ coactivator 1- α (PGC-1 α), and p53¹³.

PPAR α belongs to the nuclear receptor superfamily; it functions as a transcription factor and plays crucial roles in the regulation of varieties of metabolic dysfunctions related to inflammatory response, glucose and lipid

¹Department of Physiology, Keimyung University School of Medicine, Daegu 42601, Korea. ²Department of Internal Medicine, Keimyung University School of Medicine, Daegu 41931, Republic of Korea. ³Department of Life Sciences, Ewha Womans University, Seoul 03760, Republic of Korea. ⁴Division of Life Sciences, Korea University, 145 Anam-Ro, Seongbuk-Gu, Seoul, 02841, Republic of Korea. ⁵Department of Biochemistry, Chonbuk National University Medical School, Jeonju, Jeonbuk 54896, Republic of Korea. ⁶Division of Endocrinology and Metabolism, Department of Internal Medicine, Kyungpook National University School of Medicine, Daegu, 41944, Republic of Korea. ⁷National Creative Research Initiatives Center for Nuclear Receptor Signals, Hormone Research Center, School of Biological Sciences and Technology, Chonnam National University, Gwangju, 61186, Republic of Korea. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to S.-S.I. (email: ssim73@kmu.ac.kr)




Figure 1. Metformin increases IGFBP-2 gene expression in primary hepatocytes. DNA microarray analysis of gene expression in the metformin-treated primary hepatocytes. The ratios of gene profiles were presented by either heatmap (A) or gene expression pattern (B). (B) Representative values from the DNA microarray. Relative *Igfbp-1* and *Igfbp-2* mRNA levels with metformin treatment compared to control group in primary hepatocytes. (C) *Igfbp-2* expression in primary hepatocytes stimulated by metformin (12 h) for the indicated dose. * $p < 0.05$, ** $p < 0.01$ vs. untreated control.

metabolism¹⁴. PPAR α is expressed in various tissues including the liver, adipose tissue, heart, kidney, and intestine^{14,15}. PPAR α is activated during fasting or by ligands; PPAR α induces fatty acid oxidation and gluconeogenesis¹⁶. In addition, PPAR α is known to suppress inflammation and to preserve insulin sensitivity¹⁷. It also interacts with the retinoid X receptor, and the resulting heterodimer promotes the transcriptional activation of target genes by binding to the consensus PPAR response element (PPRE) on target gene promoters¹⁸. Further, PPAR α regulates cancer metabolism by attenuating IGF-1R signaling and Akt phosphorylation in various cancer cells^{19,20}. However, a potential link between the Sirt1-PPAR α axis and IGF-1 signaling system has not been elucidated yet.

The antidiabetic drug, metformin, is widely used for the treatment of type 2 diabetes. Metformin attenuates hepatic glucose production and triglycerides accumulation by ameliorating hyperglycemia and fatty acid oxidation in the liver^{21–23}. Metformin also improves hepatic dysfunction by stimulating liver kinase B-1, which promotes the expression of AMP-activated protein kinase (AMPK) in the liver²⁴. Because AMPK functions as a potential intracellular energy sensor and a master regulator of metabolic homeostasis, understanding the mechanisms of its activation by various physiological stimuli or therapeutic drugs and several hormones including adiponectin and leptin²⁵ are of utmost importance in developing anti-diabetic drugs.

In this study, we elucidated the potential role of IGFBP-2 in diet-induced obesity or diabetic mice. Our results demonstrated that metformin controls *Igfbp-2* gene transcription through the AMPK-Sirt1-PPAR α signaling pathway. Moreover, we showed that regulation of the Sirt1-PPAR α -IGFBP-2 signaling cascade by AMPK activator represents a novel pathway that could be applied to ameliorate metabolic syndromes by controlling IGF-1 homeostasis.

Results

Metformin increases *Igfbp-2* gene expression in primary hepatocytes. *Igfbp-2* gene expression is decreased in diabetic condition and increased when insulin resistance was improved in diabetic animal model¹⁰. We first performed DNA microarray analysis and compared global mRNA expression patterns in the primary hepatocytes by gene profiling to find putative target genes of metformin. Among genes that exhibited significant change in expression of 1.5-fold or higher, IGFBPs gene with control genes are listed in Fig. 1A. Because metformin as an anti-diabetic drug is able to improve insulin sensitivity in diabetic patients²⁶, we have speculated that metformin might be responsible for the regulation of *Igfbp-2* gene expression and biological function. The expression level of *Igfbp-2* was significantly increased in the metformin-treated primary hepatocytes (Fig. 1B). Next, to confirm microarray data under *in vitro* conditions, we performed quantitative polymerase chain reaction (qPCR) analysis on metformin-exposed mouse primary hepatocytes. As shown in Fig. 1C, *Igfbp-2* expression was markedly increased by metformin compared to untreated control ($p < 0.05$); however, *Igfbp-1* expression was not induced by metformin treatment (Fig. 1C). Together, these findings demonstrate that metformin upregulates *Igfbp-2* expression *in vitro*.

Hepatic *Igfbp-2* gene expression in HFD, ob/ob, and db/db mice. To determine the potential role of *Igfbp-2* in obese and diabetic states, we observed the expression level in mouse liver. *Igfbp-2* expression was

significantly decreased in the livers of high-fat diet-fed (HFD), *ob/ob* and *db/db* mice ($p < 0.05$); however, *Igfbp-1* was not significantly different. Lipogenic gene expressions (*Acc1* and *Gpat*) were increased in HFD, *ob/ob*, and *db/db* mice, which were measured as positive controls (Fig. 2A–C). Interestingly, the expression level of hepatic *Igfbp-1* as well as lipogenic genes was significantly increased in *db/db* mice (Fig. 2C). To observe the effects of metformin on IGFBP-2 level *in vivo*, *ob/ob* and *db/db* mice were treated by oral gavage with 100 mg/kg/day metformin for 3 weeks and observed increased IGFBP-2 level from serum and livers. Metformin administration was induced significantly an increase of *Igfbp-2* mRNA level in the livers of *ob/ob* and *db/db* mice and phosphorylation of AMPK protein was also increased in the livers of metformin-administrated lean and *db/db* mice (Fig. 2D,E). As was the case with the mRNA expression, serum levels of IGFBP-2 were also significantly increased in the metformin-fed mice, particularly in the *ob/ob* and *db/db* mouse rather than their littermates control groups (Fig. 2F). Taken together, these results suggested that *Igfbp-2* might be involved in the development obesity- and diabetes-associated metabolic dysfunction of the liver.

Metformin-induced *Igfbp-2* gene expression is mediated by AMPK. To determine whether metformin-mediated AMPK activation is involved in the induction of *Igfbp-2*, we examined the effect of metformin-AMPK pathway on IGFBP-2 after treating primary hepatocytes with metformin and compound C (Com C), an AMPK inhibitor. Com C significantly repressed metformin-mediated induction of *Igfbp-2* mRNA and protein (Fig. 3A,B). Next, to confirm that metformin-mediated *Igfbp-2* expression occurs through AMPK, we adopted adenovirus-mediated gene silencing of AMPK (Ad-si *Ampkα2*) in the AML12 cell line. As shown in Fig. 3C, Ad-si *Ampkα2* significantly decreased IGFBP-2 protein level in the AML12 cells treated with metformin. In addition, Ad-si *Sirt1* treatment also decreased metformin-induced IGFBP-2 protein level (Fig. 3C). To further demonstrate effect of AMPK α 1 on *Igfbp-2* gene regulation, mouse primary hepatocytes were infected with adenovirus overexpressing dominant negative form of AMPK (Ad-DN-*Ampk*). As expected, up-regulation of *Igfbp-2* mRNA and protein levels by metformin were markedly attenuated by Ad-DN-*Ampk* when compared with that of the controls (Supplementary Fig. 1). In addition, we measured *Ampkα1* and *Ampkα2* mRNA expression levels in the Ad-si *Sirt1* infected AML12 cell line and Ad-*Sirt1* infected mouse primary hepatocytes with metformin treatment to confirm whether *Sirt1* expression affects *Ampk* expression. Increased *Ampkα1* and *Ampkα2* mRNA expression levels by metformin were significantly decreased by *Sirt1* depletion and were increased by *Sirt1* overexpression (Supplementary Fig. 2). Next, to confirm that metformin mediates *Igfbp-2* expression through PPAR α , we overexpressed *Sirt1* using adenovirus with metformin treatment in *Pparα* null primary hepatocytes and measured *Igfbp-2* expression level. As expected, in *Pparα* null cells, even when *Sirt1* is overexpressed, metformin was unable to increase *Igfbp-2* mRNA expression level (Fig. 3D). This result suggests that *Sirt1* may be involved in regulation of *Igfbp-2* expression essentially through PPAR α . Collectively, these results indicate that AMPK plays a major role in regulating metformin-mediated stimulation of *Igfbp-2* expression in primary hepatocytes and the AML12 cell line and *Sirt1* could be also involved in the transcriptional regulation of metformin-mediated *Igfbp-2* gene expression.

***Sirt1* controls *Igfbp-2* expression via PPAR α -dependent pathway.** To further confirm the relationship between *Sirt1* and *Igfbp-2* gene expression *in vivo*, we performed qPCR analysis on liver samples obtained from obese and diabetic mice. The expression levels of *Sirt1* and *Pparα* mRNA were significantly decreased in the obese and diabetic mice compared to respective control groups (Fig. 4A). Metformin significantly elevated *Sirt1* and *Pparα* mRNA levels which were markedly diminished by Com C in primary hepatocytes (Fig. 4B). Next, we investigated whether *Sirt1* is also involved in the regulation of *Igfbp-2* expression by silencing *Sirt1* (Ad-si *Sirt1*). We infected adenovirus of si *Sirt1* in primary hepatocytes. Metformin significantly elevated *Igfbp-2* and *Pparα* mRNA levels, and this stimulatory effect of metformin was markedly attenuated by Ad-si *Sirt1* (Fig. 4C). Interestingly, the increase in the mRNA levels of *Igfbp-2* and *Pparα* was observed in metformin-administered wild-type (WT) mice but not in *Pparα*-null mice (Fig. 4D). Furthermore, metformin increased IGFBP-2 secretion in primary hepatocytes isolated from WT mice, whereas *Ampkα2*- or *Sirt1*-silenced primary hepatocytes exhibited significantly reduced IGFBP-2 secretion compared with primary hepatocytes from *Pparα*-null mice (Fig. 4E). Taken together, these results suggest that the metformin-AMPK-*Sirt1* signaling network controls *Igfbp-2* gene expression in a PPAR α -dependent manner.

Metformin regulates PPAR α -mediated *Igfbp-2* gene promoter activity. To investigate whether metformin modulates the promoter activity of mouse *Igfbp-2*, we performed transient transfection reporter assay using mouse *Igfbp-2*-Luc (*mIgfbp-2*-Luc) and *Sirt1* and *Pparα* expression vectors. As expected, the transcriptional activity of *Igfbp-2* gene promoter activity was significantly increased by metformin treatment and by overexpression of *Sirt1* or *Pparα* genes (Fig. 5A). Serial deletion and/or PPRE deletion of the *Igfbp-2* promoter was performed. Promoter activity of *Igfbp-2* gene was increased by overexpression of *Pparα* and truncations of the PPRE on the promoter resulted in a loss of PPAR α responsiveness (Fig. 5B). Moreover, metformin and PPAR α -mediated activation of *Igfbp-2* gene promoter activity was decreased by internal deletion of PPRE on full length of *Igfbp-2* gene promoter (Fig. 5C). ChIP assays of mouse primary hepatocytes in the presence of metformin showed that PPAR α occupancy was greater on the *Igfbp-2* promoter when metformin was treated (Fig. 5D). These results strongly suggest that metformin positively regulates the *Igfbp-2* promoter through PPAR α occupancy.

Metformin controls IGF-1 signaling cascade in primary cultured hepatocytes. To evaluate the potential effect of PPAR α on hepatic IGF-1 bioactivity, we treated primary hepatocytes isolated from mice which were treated with IGF-1 and metformin. Metformin decreased IGF-1-induced IGF-1R and mTOR




Figure 2. Hepatic *Igfbp-2* gene expression in HFD, *ob/ob*, and *db/db* mice and the effects of *in vivo* treatment with metformin on liver tissue IGFBP-2 expression in *ob/ob* and *db/db* mice. (A) Expression of hepatic *Igfbp-2* in response to obesity and diabetes states. Quantitative polymerase chain reaction (qPCR) analysis of total RNAs from liver of chow diet (CD) and high-fat diet (HFD) mice ($n = 5$). (B) Hepatic expression of *Igfbp-2* in liver from *ob/ob* mice ($n = 5$). (C) Relative *Igfbp-2* mRNA expression in liver tissue of *db/db* mice ($n = 5$). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. CD or Lean or WT mice. Metformin was administered to lean, *ob/ob*, *db/m+*, and *db/db* mice for 3 weeks (100 mg/kg/day). (D) Total RNA was isolated from liver tissues of the mouse and the levels of IGFBP-2 mRNA expression were quantified by quantitative real-time PCR. (E) Proteins were extracted from liver tissue of the *db/m+* and *db/db* mouse administered with water or metformin. AMPK phosphorylation were detected by immunoblotting. (F) Blood sample was collected from lean, *ob/ob*, *db/m+*, and *db/db* mouse treated with water or metformin. Secretion levels of IGFBP-2 were measured by ELISA. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. vehicle administered control.

phosphorylation in primary cultured hepatocytes (Fig. 6A,B). These findings further support our data showing that metformin regulates the IGF-1 signaling system.

Metformin improves IGF-1 bioavailability in diabetic patients. Finally, we attempted to elucidate the relationship between metformin and hormonal profile in diabetic patients whose biochemical characteristics of patients were recorded. The anthropometric measurements and biochemical characteristics of all study




Figure 3. Metformin-induced *Igfbp-2* gene expression is mediated by AMPK. (A) *Igfbp-1* and *Igfbp-2* mRNA expressions in mouse primary hepatocytes treated with metformin and compound C (Com C) for 12 h. (B) Effect of metformin and compound C (Com C) on the expression of IGF1 and AMPK in primary hepatocytes. The bar graph on the right shows the quantification of IGF-1 phosphorylation over total IGF-1. (C) Effect of metformin on IGF1 protein level when *Ampk* or *Sirt1* was knocked down. AML12 cells were infected with Ad-si *Ampk*α2 and Ad-si *Sirt1* for 36 h and then treated with metformin for 24 h. IGF1 and AMPK phosphorylation in AML12 cell line for the indicated conditions were detected by immunoblotting. * $p < 0.05$, ** $p < 0.01$ vs. untreated control and/or metformin-treated cells. (D) Relative hepatic *Igfbp-2*, *Sirt1*, and *Ppara* mRNA level in the primary hepatocytes from WT and *Ppara* null mice transfected with Ad-GFP or Ad-*Sirt1*. Mouse primary hepatocytes from WT and *Ppara* null mice were infected with Ad-GFP or Ad-*Sirt1* for 24 h and then treated with metformin for 12 h. *** $p < 0.001$ vs. WT mice.

groups are summarized in Table 1, and the hormonal profiles of patients and control individuals are listed in Table 2. Blood sugar, HbA1c, markers of liver function (alanine transaminase and aspartate aminotransferase), lipid parameters (triglycerides, cholesterol, low-density lipoprotein and non-esterified fatty acid), and circulating C-reactive protein levels were significantly elevated in diabetic patients relative to control groups. These metabolic parameters were well ameliorated in patients group who took metformin. Indeed, serum IGF1 level was markedly reduced in diabetic patients. On the contrary, circulating IGF1 level of metformin treatment group was restored to similar with that observed in control group (Fig. 7A). Interestingly, even though IGF-1 level is huge increased in diabetic patients, metformin treatment did not suppress an increased IGF-1 level in diabetic patients




Figure 4. *Sirt1* controls *Igfbp-2* expression via a *Pparα*-dependent. (A) Relative *Sirt1* and *Pparα* mRNA expression in liver tissue from HFD, *ob/ob*, and *db/db* mice ($n = 5$). $*p < 0.05$, $**p < 0.01$, $***p < 0.001$ vs. CD or Lean or WT mice. (B) Expression of hepatic *Sirt1* and *Pparα* in mouse primary hepatocytes treated with metformin and compound C (Com C) for 12 h. (C) qPCR analysis of hepatic *Sirt1*, *Igfbp-2*, and *Pparα* mRNA level in the primary hepatocytes transfected with Ad-si *Sirt1*. Mouse primary hepatocytes were transfected with Ad-si *Sirt1* for 36 h and then treated with metformin for 12 h. (D) Mouse primary hepatocytes from WT and *Pparα* null mice were incubated in the presence or absence of metformin for 12 h *Igfbp-2* and *Pparα* mRNA expression were measured by qPCR analysis. (E) Secretion level of IGFBP-2 *in vitro*. Mouse primary hepatocytes from WT and *Pparα* null mice were infected with Ad-si *Ampkα2* and Ad-si *Sirt1* for 36 h and then treated with metformin for 24 h. Supernatant medium were collected and secretion levels of IGFBP-2 were measured by ELISA. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$ vs. untreated control or metformin-treated cells.




Figure 5. Metformin regulates PPAR α -mediated *Igfbp-2* gene expression. (A) Effect of the *Ppar α* on mouse *Igfbp-2* promoter activity. Transient transfections were performed using observed promoter construct, *Igfbp-2* (–1314/+78) in 293T cell. Below, immunoblot analysis of *Sirt1* and *Ppar α* expression in transfected HEK293T cells, detected with anti-*Sirt1* and PPAR α . Equal amounts of protein were confirmed using β -actin. (B) Schematic drawing of *Ppar α* binding region of *Igfbp-2* promoter. Serial deletion constructs of *Igfbp-2* promoter (–698/–686 bp) were transiently transfected with *Ppar α* to 293T cell. (C) PPRE-dependent activation of *Igfbp-2* promoter activity. Alignment of potential PPRE in *Igfbp-2* promoter is indicated (top panel). (D) ChIP assay. Mouse primary hepatocytes were treated metformin for 24 h. Chromatin was precipitated using an anti-PPAR α antibody from primary hepatocytes, and purified DNA samples were used to perform qPCR with primers binding to the PPRE regions on the *Igfbp-2* gene promoter. Input represents 10% of purified DNA in each sample. * $p < 0.05$ vs. untreated control or individual-treated cells. mlgfbp-2, mouse *Igfbp-2*.

(Fig. 7B). These findings suggest that metformin plays a pivotal role in regulating IGF-1 bioavailability without reducing total IGF-1 level in diabetic patients.

Discussion

In this study, we demonstrate that metformin increases *Igfbp-2* expression through the AMPK-Sirt1-PPAR α signaling cascade both *in vitro* and *in vivo*. Conversely, these effects of metformin were attenuated by si *Ampk α 2* over-expression as well as silencing of *Sirt1* and *Ppar α* genes. Further analysis revealed that metformin stimulates the *Igfbp-2* promoter activity by PPAR α . Previously we have reported the presence of PPAR α response element on *Igfbp-2* promoter that contributes to fasting induction of IGFBP-2²⁷. Based on these data, we speculated that metformin decreased IGF-1 action by upregulating *Igfbp-2* gene expression through an AMPK-Sirt1-PPAR α signaling network.

Circulating IGFBP-2 levels are correlated with metabolic dysfunction including obesity, diabetes, and insulin resistance as *Igfbp-2* gene expression is lower in obese and diabetic patients²⁸. IGFBP-2 is a negative regulator of IGF-1 homeostasis^{7,8,29}. A previous study showed IGFBP-2 concentrations were lower in serum of mildly insulin-resistant glucose-tolerant patients²⁸. Also, restoration of IGFBP-2 in insulin-resistant patients is associated with improved insulin sensitivity^{30,31}. These clinical observations are consistent with the results of our study, where a decrease in *Igfbp-2* expression and serum levels were observed in both diet-induced obese and diabetic




Figure 6. Metformin controls IGF-1 signaling cascade in primary cultured hepatocytes. (A) Mouse primary hepatocytes were pretreated with metformin for 12 h and then exposed to IGF-1 for 15 min at the indicated conditions. Whole cell extracts were isolated from primary hepatocytes of the indicated groups and measured by western blot analysis with various antibodies. (B) The left panel reveals the quantification of IGF-1 phosphorylation over total IGF-1, right panel indicates mTOR phosphorylation over total mTOR, and bottom panel indicates p70S6K phosphorylation over total p70S6K. * $p < 0.05$, ** $p < 0.01$ vs. untreated control or IGF-1-treated cells.

Characteristic	Nor	DM	DM + Met
N(M/F)	53 (17/36)	20 (9/11)	36 (21/15)
Age (years)	58.33 ± 3.83	54.31 ± 3.36	63.53.31 ± 2.95
Height (cm)	158.81 ± 1.95	163.74 ± 1.77	159.92 ± 2.12
Weight (g)	62.02 ± 2.26	67.62 ± 3.38	65.13 ± 3.15
BMI (kg/m ²) (male)	24.03 ± 3.03	26.16 ± 0.89	25.6 ± 3.13
BMI (kg/m ²) (female)	24.96 ± 2.84	23.24 ± 1.56	24.88 ± 4.68

Table 1. Clinical characteristics of the participants.

Characteristic	Nor	DM		DM + Met	
FBS (mg/dl)	94.58 ± 5.16	132.92 ± 13.51	**	117.53 ± 7.88	**
HbA1c (mmol/l)	6.18 ± 0.13	8.16 ± 0.62	*	6.85 ± 0.15	*
Insulin (pmol/l)	32.56 ± 4.97	24.08 ± 4.38	**	24.73 ± 2.66	
C-peptide	10.4 ± 0.87	8.03 ± 1.1	*	7.9 ± 0.97	
ALT (IU/l)	26.17 ± 5.27	28.69 ± 4.44	*	19.8 ± 2.51	**
AST (IU/l)	30.83 ± 2.8	25.23 ± 3.38	*	20.4 ± 1.17	*
TG (mg/dL)	147.36 ± 38.4	152.22 ± 17.2	*	118.69 ± 14.64	**
Chol (mg/dl)	165.75 ± 11.06	193.38 ± 12.43	**	164.53 ± 11.22	**
HDL (mg/dl)	48.43 ± 3.01	44.73 ± 3.88	*	43.87 ± 1.74	
LDL (mg/dl)	96.35 ± 9.88	113.28 ± 10.87	*	101.87 ± 9.2	**
CRP (mg/dl)	3.03 ± 1.31	9.59 ± 6.51	***	1.39 ± 0.51	***
NEFA (μEg/l)	762.08 ± 125.83	562.31 ± 80.32	**	522.13 ± 78.28	*

Table 2. Biochemical characteristics of the participants with metformin-challenged type2 diabetes mellitus. Nor-Normal; DM-Diabetes mellitus; Met-Metformin. FBS-fasting blood sugar; ALT-Alanine aminotransferase; AST-Aspartate aminotransferase. TG-Triglyceride; Chol-Cholesterol; HDL-High-density lipoprotein; LDL-Low Density lipoprotein; CRP-C-reactive protein; NEFA-Non-esterified fatty acids. Data presented as Mean ± SEM, n = 12–15, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ vs. Nor or DM.




Figure 7. Metformin improves IGF-1 bioavailability in diabetic patients. Serum IGFBP-2 (A) and IGF-1 (B) in the control normal, diabetes (DM), and metformin-treated diabetes patients (DM + Met). (C) Scheme of the role of metformin in the regulation of IGFBP-2-IGF-1 signaling network. Metformin elevates *Igfbp-2* gene expression through AMPK-Sirt1-PPAR α signaling pathway and then improves IGF-1 bioavailability via attenuation of IGF-1 signaling. However, the beneficial effects of metformin are counteracted in diet-induced obesity and diabetes states.

animal models as well as in diabetic patients compared to control groups. IGF-1 levels were also significantly higher in diabetic patients. From these findings, we propose that IGFBP-2 may modulate hepatic metabolic dysfunction by regulating IGF-1 homeostasis.

As mentioned previously, *Sirt1* is associated with metabolic diseases involving the IGF-1 signaling system in the liver^{12,32,33}. However, the connection between the IGFBP-2-IGF-1 axis and the beneficial effects of *Sirt1* in the liver was not previously understood. We demonstrated that metformin significantly elevated hepatic *Sirt1* gene expression in an AMPK-dependent manner. We verified that metformin-induced *Igfbp-2* expression was mediated by the Sirt1-dependent pathway and the effect of metformin was disrupted by gene silencing of *Sirt1*. Thus, our studies strongly suggest that regulation of the metformin-AMPK pathway through Sirt1 plays an important role in ameliorating hepatic metabolic dysfunction by controlling *Igfbp-2* gene expression. However, there is a report suggesting that Sirt1 affects AMPK activation³⁴. In our data, we showed that metformin-induced *Ampk α 1* and *Ampk α 2* mRNA expression levels were decreased by knockdown of *Sirt1* and were increased by *Sirt1* overexpression (Supplementary Fig. 2). These results indicated that AMPK expression might be also regulated by Sirt1 in a feedback regulatory mechanism.

PPAR α acts a key role in regulating diverse metabolic processes and maintaining metabolic balance in response to changes in fuel availability^{14,15}. A recent study demonstrated that AMPK activators such as metformin and panduratin A control diet-induced obesity and inflammation through the PPAR α pathway in the liver³⁵. Therefore, it has been suggested that AMPK activators, including resveratrol, EGCG, and other natural products, might have beneficial effects on hepatic metabolic dysfunction. We also demonstrated that the metformin-AMPK-PPAR α pathway significantly activates *Igfbp-2* expression in mice and that metformin treatment is correlated with increased serum levels of IGFBP-2 in human patients.

This report suggests that metformin may be involved in IGF-1 homeostasis in response to metabolic dysfunction, and this effect was disrupted by gene silencing of AMPK expression and in *Ppar α* -null mice. Our results imply that activation of AMPK and PPAR α in human patients might be possible to regulate the IGFBP-2-IGF-1 axis. Meanwhile, there is possibility that another signaling pathway might exist between the IGFBP-2-IGF-1 axis and the metformin-AMPK-PPAR α pathway in the liver. Moreover, interestingly, we showed *Ppar α* expression level was increased by metformin treatment in primary hepatocytes and AML12 cells. Indeed there were putative transcription factors binding sites on the *Ppar α* promoter. Even though we did not provide a mechanistic mechanism for how metformin actually induced *Ppar α* gene expression here, the signaling system underlying these effects of metformin needs to be elucidated in future studies. Also, even though IGFBP-2 secretion was completely inhibited, metformin treatment of PPAR-null cell showed slight induction of *Igfbp-2* expression (Fig. 4D). We speculated that transcription factors other than PPAR α may be involved in the regulation of *Igfbp-2* gene expression. Moreover, IGFBP-2 secretion has been known to be stimulated by growth hormone³⁶, but the underlying regulatory mechanism has not been addressed yet. Understanding the discrepancy between intracellular and secretory IGFBP-2 regulation could be an interesting research in the future.

In conclusion, as depicted in the novel schematic model in Fig. 7C, our current study suggests that upregulation of *Igfbp-2* by metformin activates the AMPK-Sirt1-PPAR α signaling network. Furthermore, our results also indicate that metformin affects IGF-1 homeostasis by modulating IGF-1 bioavailability in the diabetic condition. These findings may aid the development of a novel pathway that could be applied to ameliorate metabolic dysfunction.

Materials and Methods

Reagents. Metformin (Sigma-Aldrich, St. Louis, MO, USA), compound C (Calbiochem, Billerica, MA, USA), and recombinant human IGF-1 (Life Technologies, Carlsbad, CA, USA) were purchased from the indicated companies and dissolved in the recommended solvents. Cell culture media were purchased from Gibco-BRL (Grand Island, NY, USA).

Animal studies. All animal experiments were performed in accordance with the approved guidelines and the protocol is approved by the Institutional Animal Use and Care Committee (IAUCC), Keimyung University School of Medicine (KM-2014-33R3). Male C57BL6 mice, *ob/ob*, *db/db* (Jung-Ang Experimental animals, Seoul, Republic of Korea), and *Ppar α* null mice (8-weeks-of-age) were used in experiments, as previously described³⁷. For the diet experiment, 8-week-old male mice were fed with a LFD (10% of total calories in the form of lard fat) or HFD (60% of total calories in the form of lard fat) for 12 wks (D12450B or D12492; Research Diets, New Brunswick, NJ, USA).

Metformin administration. The male mouse (7 weeks, $n = 32$) strains *db/m+* (C57BLKS/J *lar-m⁺/Lepr^{db}*), *db/db* (C57BLKS/J *lar-Lepr^{db}/Lepr^{db}*), lean and *ob/ob* (C57BL/6J Ham Slc-*ob/ob*) were divided into two groups: metformin ($n = 16$) and control groups ($n = 16$). In the metformin group, metformin (100 mg/kg/day) was administered by the oral for 3 weeks. In control group, vehicle (water) was administered by daily oral gavage. After the mice were fasted for 12 h at the end of the treatment period, blood samples were collected. Blood samples were used for determinations of serum IGFBP-2 levels. Real-time PCR and Western blot analysis were performed to examine IGFBP-2 mRNA and protein expression in the removed liver tissues, respectively. All animal experiments were carried out in accordance with the institutional guidelines and the all procedures and care administered were approved by the Institutional Animal Use and Care Committee (IAUCC), Keimyung University School of Medicine (KM-2012-55R and KM-2016-3R).

Construction of plasmids and DNA. The cDNA encompassing the *Igfbp-2* open reading frame was cloned by reverse transcriptase-PCR. Primers containing *EcoRI* were used for cloning of mouse *Igfbp-2*. The PCR product was digested by *EcoRI* and ligated into pcDNA3 (Invitrogen, CA, USA). The reporter plasmid mouse *Igfbp-2* promoter was PCR-amplified from mouse genomic DNA (Novagen, Merck KGaA, Darmstadt, Germany) and inserted into the pGL3 Basic vector (Promega, Madison, WI, USA) using the *SmaI* and *EcoRI* restriction enzyme sites. Expression vector for *Sirt1* and *Ppar α* were previously described³⁸. The deletion form of mouse *Igfbp-2*-Luc was generated using PCR analysis. Plasmids were confirmed by sequencing analysis.

Cell culture and transient transfection assays. AML-12 immortalized mouse hepatocytes were cultured in DMEM/F-12 medium (Gibco-BRL) supplemented with 10% FBS, insulin-transferrin-selenium (Gibco-BRL), dexamethasone (40 ng/mL; Sigma-Aldrich), and antibiotics in a humidified atmosphere containing 5% CO₂ at 37 °C. HEK293T cells were cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco-BRL, Grand Island, NY, USA) supplemented with 10% fetal bovine serum (FBS; Hyclone, Logan, UT, USA) and antibiotics in a humidified atmosphere containing 5% CO₂ at 37 °C. Transient transfections were conducted using Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instruction.

Isolation and culture of primary mouse hepatocytes. Mouse primary hepatocytes were isolated from the livers of 8-week-old male mice (Samtako, Osan, Republic of Korea). The protocol for isolation of hepatocytes was described previously³⁹.

Microarray expression profiling and CyberT analysis. Total RNA was prepared from each primary hepatocytes separately and equal amounts from 9 different individual samples were pooled together. Equal amounts of RNA (500 ng) from each pool were used to hybridize to triplicate microarray slides covering the entire

transcriptome (Operon Whole 27K Oligo chip from Operon). The results for the triplicate chips were analyzed by GenePix operating software (GenePix pro 4.1). For statistical and biological significance, we set the Standard Deviation of Differential Expression ($SD < p$) at <0.5 and fold change at 1.5. These data were used to construct the “heat-map” using MeV v4.61 multiexperiment viewer software (<http://www.tm4.org/mev>).

Recombinant adenovirus. Adenoviruses expressing US, sh *Ampk α 2*, si *Sirt1*, *Sirt1*, and green fluorescence protein (GFP) have been described previously^{40,41}.

Measurement of mRNA. Total RNA was isolated from liver tissue and used for qPCR as previously described³⁷. The expression of all transcripts using qPCR data were normalized to ribosomal L32 expression. The sequences of the primers used for gene expression analyses are reported in Supplementary Table 1.

Immunoblotting. Proteins were isolated from liver tissues and analyzed according to the methods described previously. The membranes were probed with antibody against phospho-AMPK, AMPK, phospho-IGF-1R, phospho-Akt, Akt, phospho-Sirt1, Sirt1, phospho-p70S6K, p70S6K (Cell Signaling Technology, Danvers, MA, USA), PPAR α , IGFBP-2, IGF-1R, and β -actin (Santa Cruz Biotechnology, Santa Cruz, CA, USA), and then developed using an ECL Western blot detection kit (Amersham Bioscience, Piscataway, NJ, USA).

Human study population. This study was carried out in adherence with the guidelines of the Declaration of Helsinki and approved by the Institutional Review Boards of Keimyung University Dongsan medical center in Korea (DSMC2013-09-003-010), and informed consents were obtained from all subjects. The present study was conducted on patients with type 2 diabetes ($n = 56$) and non-diabetic control ($n = 53$) who were confirmed with insulin secretion test in the department of Internal Medicine in Dongsan medical center from October 2012 to October 2014. Among the 56 patients with type 2 diabetes, the prevalence of metformin medication was 64.3% ($n = 36$) and medication without metformin was 35.7% ($n = 20$). Exclusion criteria were: chronic alcohol drinking, hepato-biliary abnormalities, or any other acute disease.

Human study design and assessments. A detailed questionnaire was completed for each of the 109 participating subjects. Information obtained included age, gender, height, weight, blood pressure, smoking history, history of alcohol consumption or hepatobiliary disorders, duration of diabetes, and history of hypertension or cardiovascular diseases. Additionally, fasting blood glucose (FBG), 2 hour postprandial blood glucose (2PPBG), C-peptide, insulin, insulin-like growth factors-binding protein-2 (IGFBP-2) levels were obtained. Fasting venous blood was drawn on the day of examination via a venipuncture from the subject's antecubital vein and baseline biochemical profiles including serum total glycosylated hemoglobin (HbA1c) and glucose were analyzed using automated glycohemoglobin analyzer HLC-723G7 (Tosoh, Japan) and BIOSEN C-line, clinic (EKF Diagnostic, Germany). For serum and plasma IGFBP-2 measurement, blood samples were collected in tubes containing EDTA and IGFBP-2 concentrations were analyzed with RayBio Mouse IGFBP-2 ELISA Kit (RayBiotech, Inc., Norcross, GA, USA). This study was approved by the Institutional Review Boards of Keimyung University Dongsan medical center in Korea (DSMC2013-09-003-010), and informed consents were obtained from all subjects.

Statistical analysis. Data calculation and statistical analyses were performed using GraphPad Prism 3–5.0 software. The statistical significance of differences between groups was determined using Student's *t* test and multiple comparisons were analyzed using one-way ANOVA under treatment and experiment as factors. Results are presented as mean \pm standard deviation of at least three separate experiments. All *P* values less than 0.05 were considered significant.

References

- Oh, Y., Muller, H. L., Lee, D. Y., Fielder, P. J. & Rosenfeld, R. G. Characterization of the affinities of insulin-like growth factor (IGF)-binding proteins 1–4 for IGF-I, IGF-II, IGF-I/insulin hybrid, and IGF-I analogs. *Endocrinology* **132**, 1337–1344 (1993).
- Firth, S. M. & Baxter, R. C. Cellular actions of the insulin-like growth factor binding proteins. *Endocrine reviews* **23**, 824–854 (2002).
- Lappas, M. Insulin-like growth factor-binding protein 1 and 7 concentrations are lower in obese pregnant women, women with gestational diabetes and their fetuses. *J Perinatol* **35**, 32–38 (2015).
- Yamanaka, Y., Wilson, E. M., Rosenfeld, R. G. & Oh, Y. Inhibition of insulin receptor activation by insulin-like growth factor binding proteins. *The Journal of biological chemistry* **272**, 30729–30734 (1997).
- Arnaldez, F. I. & Helman, L. J. Targeting the insulin growth factor receptor 1. *Hematology/oncology clinics of North America* **26**, 527–542, vii–viii (2012).
- Ricort, J. M. Insulin-like growth factor binding protein (IGFBP) signalling. *Growth Horm IGF Res* **14**, 277–286 (2004).
- Hedbacker, K. *et al.* Antidiabetic effects of IGFBP2, a leptin-regulated gene. *Cell metabolism* **11**, 11–22 (2010).
- Rajaram, S., Baylink, D. J. & Mohan, S. Insulin-like growth factor-binding proteins in serum and other biological fluids: regulation and functions. *Endocrine reviews* **18**, 801–831 (1997).
- Wheatcroft, S. B. & Kearney, M. T. IGF-dependent and IGF-independent actions of IGF-binding protein-1 and -2: implications for metabolic homeostasis. *Trends in endocrinology and metabolism: TEM* **20**, 153–162 (2009).
- Laughlin, G. A., Barrett-Connor, E., Criqui, M. H. & Krititz-Silverstein, D. The prospective association of serum insulin-like growth factor I (IGF-I) and IGF-binding protein-1 levels with all cause and cardiovascular disease mortality in older adults: the Rancho Bernardo Study. *J Clin Endocrinol Metab* **89**, 114–120 (2004).
- Heald, A. H. *et al.* Insulin-like growth factor binding protein-2 (IGFBP-2) is a marker for the metabolic syndrome. *Exp Clin Endocrinol Diabetes* **114**, 371–376 (2006).
- Chang, H. C. & Guarente, L. SIRT1 and other sirtuins in metabolism. *Trends in endocrinology and metabolism: TEM* **25**, 138–145 (2014).
- Schug, T. T. & Li, X. Sirtuin 1 in lipid metabolism and obesity. *Annals of medicine* **43**, 198–211 (2011).
- Contreras, A. V., Torres, N. & Tovar, A. R. PPAR-alpha as a key nutritional and environmental sensor for metabolic adaptation. *Advances in nutrition* **4**, 439–452 (2013).

15. Monsalve, F. A., Pyarasani, R. D., Delgado-Lopez, F. & Moore-Carrasco, R. Peroxisome proliferator-activated receptor targets for the treatment of metabolic diseases. *Mediators of inflammation* **2013**, 549627 (2013).
16. Peeters, A. & Baes, M. Role of PPARalpha in Hepatic Carbohydrate Metabolism. *PPAR Res* **2010** (2010).
17. Eder, K. & Ringseis, R. The role of peroxisome proliferator-activated receptor alpha in transcriptional regulation of novel organic cation transporters. *Eur J Pharmacol* **628**, 1–5 (2010).
18. Yoon, M. The role of PPARalpha in lipid metabolism and obesity: focusing on the effects of estrogen on PPARalpha actions. *Pharmacological research: the official journal of the Italian Pharmacological Society* **60**, 151–159 (2009).
19. Drukala, J. *et al.* ROS accumulation and IGF-IR inhibition contribute to fenofibrate/PPARalpha-mediated inhibition of glioma cell motility *in vitro*. *Molecular cancer* **9**, 159 (2010).
20. Urbanska, K. *et al.* Activation of PPARalpha inhibits IGF-I-mediated growth and survival responses in medulloblastoma cell lines. *International journal of cancer. Journal international du cancer* **123**, 1015–1024 (2008).
21. Lin, H. Z. *et al.* Metformin reverses fatty liver disease in obese, leptin-deficient mice. *Nature medicine* **6**, 998–1003 (2000).
22. Kim, M. J. *et al.* Metformin-associated lactic acidosis: predisposing factors and outcome. *Endocrinol Metab (Seoul)* **30**, 78–83 (2015).
23. Zhou, G. *et al.* Role of AMP-activated protein kinase in mechanism of metformin action. *J Clin Invest* **108**, 1167–1174 (2001).
24. Shaw, R. J. *et al.* The kinase LKB1 mediates glucose homeostasis in liver and therapeutic effects of metformin. *Science* **310**, 1642–1646 (2005).
25. Zhang, B. B., Zhou, G. & Li, C. AMPK: an emerging drug target for diabetes and the metabolic syndrome. *Cell metabolism* **9**, 407–416 (2009).
26. Velazquez, E. M., Mendoza, S., Hamer, T., Sosa, F. & Glueck, C. J. Metformin therapy in polycystic ovary syndrome reduces hyperinsulinemia, insulin resistance, hyperandrogenemia, and systolic blood pressure, while facilitating normal menses and pregnancy. *Metabolism* **43**, 647–654 (1994).
27. Kang, H. S. *et al.* Regulation of IGFBP-2 expression during fasting. *The Biochemical journal* **467**, 453–460 (2015).
28. Frystyk, J., Skjaerbaek, C., Vestbo, E., Fisker, S. & Orskov, H. Circulating levels of free insulin-like growth factors in obese subjects: the impact of type 2 diabetes. *Diabetes Metab Res Rev* **15**, 314–322 (1999).
29. Wheatcroft, S. B. *et al.* IGF-binding protein-2 protects against the development of obesity and insulin resistance. *Diabetes* **56**, 285–294 (2007).
30. Rasmussen, M. H., Juul, A., Kjems, L. L. & Hilsted, J. Effects of short-term caloric restriction on circulating free IGF-I, acid-labile subunit, IGF-binding proteins (IGFBPs)-1–4, and IGFBPs-1–3 protease activity in obese subjects. *Eur J Endocrinol* **155**, 575–581 (2006).
31. Arafat, A. M. *et al.* The role of insulin-like growth factor (IGF) binding protein-2 in the insulin-mediated decrease in IGF-I bioactivity. *J Clin Endocrinol Metab* **94**, 5093–5101 (2009).
32. Yamamoto, M. *et al.* SIRT1 regulates adaptive response of the growth hormone–insulin-like growth factor-I axis under fasting conditions in liver. *Proc Natl Acad Sci USA* **110**, 14948–14953 (2013).
33. Lemieux, M. E. *et al.* The Sirt1 deacetylase modulates the insulin-like growth factor signaling pathway in mammals. *Mechanisms of ageing and development* **126**, 1097–1105 (2005).
34. Hou, X. *et al.* SIRT1 regulates hepatocyte lipid metabolism through activating AMP-activated protein kinase. *J Biol Chem* **283**, 20015–20026 (2008).
35. Buler, M. *et al.* Energy-sensing factors coactivator peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1alpha) and AMP-activated protein kinase control expression of inflammatory mediators in liver: induction of interleukin 1 receptor antagonist. *J Biol Chem* **287**, 1847–1860 (2012).
36. Pollak, M. Insulin and insulin-like growth factor signalling in neoplasia. *Nat Rev Cancer* **8**, 915–928 (2008).
37. Im, S. S. *et al.* Peroxisome proliferator-activated receptor {alpha} is responsible for the up-regulation of hepatic glucose-6-phosphatase gene expression in fasting and db/db Mice. *J Biol Chem* **286**, 1157–1164 (2011).
38. Chanda, D., Xie, Y. B. & Choi, H. S. Transcriptional corepressor SHP recruits SIRT1 histone deacetylase to inhibit LRH-1 transactivation. *Nucleic acids research* **38**, 4607–4619 (2010).
39. Im, S. S. *et al.* Glucose-stimulated upregulation of GLUT2 gene is mediated by sterol response element-binding protein-1c in the hepatocytes. *Diabetes* **54**, 1684–1691 (2005).
40. Lee, J. H. *et al.* Overexpression of SIRT1 protects pancreatic beta-cells against cytokine toxicity by suppressing the nuclear factor-kappaB signaling pathway. *Diabetes* **58**, 344–351 (2009).
41. Lee, H. M. *et al.* Acromegaly with Normal Insulin-Like Growth Factor-1 Levels and Congestive Heart Failure as the First Clinical Manifestation. *Endocrinol Metab (Seoul)* **30**, 395–401 (2014).

Acknowledgements

We thank Dr. Timothy F. Osborne (Sanford-Burnham-Prebys Medical Discovery Institute, FL, USA) and Dr. Yong-Ho Ahn (Yonsei University College of Medicine) for helpful discussion and reading critically. This study was supported by a grant of the Korea Health Technology R&D Project, Ministry of Health & Welfare, Republic of Korea (No. A120864) and supported by the National Research Foundation of Korea (NRF) Grant funded by the Korean Government (MSIP) (No. 2014R1A5A2010008) to S.S.I. This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (No. 2015R1D1A1A01057610) to H.S.K.

Author Contributions

H.S.K. and H.C.C. contributed to the design and performance of experiments, analysis and interpretation of data, writing the draft; J.H.L., G.T.O., S.H.K., B.H.P., I.K.L. and H.S.C. contributed to the performance of experiments, analysis of experimental results; H.S.C. and D.K.S. contributed to the analysis and interpretation of data, and critical review of the manuscript; S.S.I. contributed to the conception and design of the experiments, wrote the manuscript and critical review of the manuscript. S.S.I. is the guarantor of this work, as such, had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Kang, H. S. *et al.* Metformin stimulates IGFBP-2 gene expression through PPARalpha in diabetic states. *Sci. Rep.* **6**, 23665; doi: 10.1038/srep23665 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>